

Fuzzy Prime Ideals of ADL's

Chigurupalli Santhi Sundar Raj, Natnael Teshale Amare and Uppasetti Madana Swamy

Abstract—In this paper the concept of prime L -fuzzy ideals and L -fuzzy prime ideals of an ADL A with truth values in a complete lattice L satisfying the infinite meet distributive law are introduced. All prime L -fuzzy ideals of a given ADL A are determined by establishing a one-to-one correspondence between prime L -fuzzy ideals of an ADL A and the pairs (P, α) , where P is a prime ideal of A and α is a prime element in L . Also, here minimal prime L -fuzzy ideals and L -fuzzy minimal prime ideals of an ADL A are introduced and characterized.

Index Terms—Almost Distributive Lattice (ADL), complete lattice, L -fuzzy minimal prime ideal L -fuzzy prime ideal, minimal prime L -fuzzy ideal, prime L -fuzzy ideal.

I. INTRODUCTION

A fuzzy subset of a set X is a function from X into $I = [0, 1]$, as in [1]. J.A. Goguen [2] explored, generalized and continued the work of L.A. Zadeh and realized that the unit interval $[0, 1]$ is not sufficient to take the truth values of general fuzzy statements. Wang-Jing Liu [3] introduced the notion of a fuzzy ideal of a ring in the case when $L = [0, 1]$ of real numbers and T.K. Mukherjee and M.K. Sen [4] introduced the notion of a fuzzy prime ideal and continued the study of fuzzy ideals. U.M. Swamy and K.L.N. Swamy [5] introduced the concept of fuzzy prime ideal of a ring with truth values in a complete lattice satisfying the infinite meet distributive law.

The concept of prime ideal of an Almost Distributive Lattice was introduced by U.M. Swamy and G.C. Rao, in 1981 [6]. U.M. Swamy, Ch. Santhi Sundar Raj and Natnael Teshale A [7] have introduced the notion of L -fuzzy ideals of an ADL with the truth values in a complete lattice L satisfying the infinite meet distributive law.

In this paper, we introduce and study prime L -fuzzy ideals and L -fuzzy prime ideals of an ADL A , where L is a complete lattice satisfying the infinite meet distributive law. Also, in this paper we introduce minimal prime L -fuzzy ideals and L -fuzzy minimal prime ideals of an ADL A .

II. PRELIMINARIES

First we give necessary definitions and results mostly taken from [6] and [7] which will be used in the later text.

Definition 2.1: An algebra $A = (A, \wedge, \vee, 0)$ of type $(2, 2, 0)$ is called an Almost Distributive Lattice (abbreviated as ADL) if it satisfies the following conditions for all a, b and $c \in A$.

- 1) $0 \wedge a = 0$

- 2) $a \vee 0 = a$
- 3) $a \wedge (b \vee c) = (a \wedge b) \vee (a \wedge c)$
- 4) $a \vee (b \wedge c) = (a \vee b) \wedge (a \vee c)$
- 5) $(a \vee b) \wedge c = (a \wedge c) \vee (b \wedge c)$
- 6) $(a \vee b) \wedge b = b$.

Any bounded below distributive lattice is an ADL, where 0 is the smallest element. Any nonempty set X can be made into an ADL by fixing an arbitrarily chosen element 0 in X and by defining the binary operations \wedge and \vee on X by

$$a \wedge b = \begin{cases} 0, & \text{if } a = 0 \\ b, & \text{if } a \neq 0 \end{cases} \quad \text{and} \quad a \vee b = \begin{cases} b, & \text{if } a = 0 \\ a, & \text{if } a \neq 0. \end{cases}$$

This ADL $(X, \wedge, \vee, 0)$ is called a discrete ADL.

Definition 2.2: Let $A = (A, \wedge, \vee, 0)$ be an ADL. For any a and $b \in A$, define $a \leq b$ if $a = a \wedge b$ ($\Leftrightarrow a \vee b = b$). Then \leq is a partial order on A with respect to which 0 is the smallest element in A .

Theorem 2.3: The following hold for any a, b and c in an ADL A .

- (1) $a \wedge 0 = 0 = 0 \wedge a$ and $a \vee 0 = a = 0 \vee a$
- (2) $a \wedge a = a = a \vee a$
- (3) $a \wedge b \leq b \leq b \vee a$
- (4) $a \wedge b = a \Leftrightarrow a \vee b = b$
- (5) $a \wedge b = b \Leftrightarrow a \vee b = a$
- (6) $(a \wedge b) \wedge c = a \wedge (b \wedge c)$ (i.e., \wedge is associative)
- (7) $a \vee (b \vee a) = a \vee b$
- (8) $a \leq b \Rightarrow a \wedge b = a = b \wedge a$ ($\Leftrightarrow a \vee b = b = b \vee a$)
- (9) $(a \wedge b) \wedge c = (b \wedge a) \wedge c$
- (10) $(a \vee b) \wedge c = (b \vee a) \wedge c$
- (11) $a \wedge b = b \wedge a \Leftrightarrow a \vee b = b \vee a$
- (12) $a \wedge b = \inf\{a, b\} \Leftrightarrow a \wedge b = b \wedge a \Leftrightarrow a \vee b = \sup\{a, b\}$.

An element $m \in A$ is said to be maximal if, for any $x \in A$, $m \leq x$ implies $m = x$. It can be easily observed that m is maximal if and only if $m \wedge x = x$ for all $x \in A$.

Definition 2.4: Let I be a non empty subset of an ADL A . Then I is called an ideal of A if $a, b \in I \Rightarrow a \vee b \in I$ and $a \wedge x \in I$ for all $x \in A$.

As a consequence, for any ideal I of A , $x \wedge a \in I$ for all $a \in I$ and $x \in A$. For any $S \subseteq A$, the smallest ideal of A containing S is called the ideal generated by S in A and is denoted by $[S]$. It is known that

$$[S] = \left\{ \left(\bigvee_{i=1}^n x_i \right) \wedge a \mid n \geq 0, x_i \in S \text{ and } a \in A \right\}.$$

when $S = \{x\}$, we write $[x]$ for $\{[x]\}$. Note that $[x] = \{x \wedge a \mid a \in A\}$.

Definition 2.5: An L -fuzzy subset λ of X is a mapping from X into L , where L is a complete lattice satisfying the infinite meet distributive law. If L is the unit interval $[0, 1]$ of real numbers, then these are the usual fuzzy subsets of X .

Manuscript received December 19, 2017; accepted May 10, 2018.

Ch. Santhi Sundar Raj and A. Natnael Teshale are with the Department of Engineering Mathematics, Andhra University, Visakhapatnam- 530003, A.P., India. E-mail: santhisundarraaj@yahoo.com

A. Natnael Teshale is with the Department of Mathematics, University of Gondar, Ethiopia. E-mail: yenaatnaelau@yahoo.com

U.M. Swamy is with the Department of Mathematics, Andhra University, Visakhapatnam- 530003, A.P., India. E-mail: umswamy@yahoo.com

For any $\alpha \in L$, the set $\lambda_\alpha = \{x \in X : \alpha \leq \lambda(x)\}$ is called the α -cut of λ .

Definition 2.6: An L -fuzzy subset λ of A is said to be an L -fuzzy ideal of A , if $\lambda(0) = 1$ and $\lambda(x \vee y) = \lambda(x) \wedge \lambda(y)$, for all $x, y \in A$.

Lemma 2.7: Let λ be an L -fuzzy ideal of A , S a non-empty subset of A and $x, y \in A$. Then we have the following.

- (1) $x \wedge y = y$ and $y \wedge x = x \implies \lambda(x) = \lambda(y)$
- (2) $\lambda(x \wedge y) = \lambda(y \wedge x)$
- (3) $x \in [S] \implies \lambda(x) \geq \bigwedge_{i=1}^n \lambda(a_i)$ for some $a_1, a_2, \dots, a_n \in S$
- (4) $x \in [y] \implies \lambda(x) \geq \lambda(y)$
- (5) If m is a maximal element in A then $\lambda(m) \leq \lambda(x)$, for all x
- (6) $\lambda(m) = \lambda(n)$ for all maximal elements m and n in A .

Theorem 2.8: The set of all L -fuzzy ideals of A is a complete distributive lattice, in which the supremum $\bigvee_{i \in \Delta} \lambda_i$ and infimum $\bigwedge_{i \in \Delta} \lambda_i$ of any family $\{\lambda_i : i \in \Delta\}$ of L -fuzzy ideals of A are given by

$$\left(\bigvee_{i \in \Delta} \lambda_i\right)(x) = \bigvee_{a \in F} \left\{ \bigwedge_{i \in \Delta} \left(\bigvee_{i \in \Delta} \lambda_i(a)\right) : x \in (F], F \subset \subset A \right\}$$

$$\text{and } \left(\bigwedge_{i \in \Delta} \lambda_i\right)(x) = \bigwedge_{i \in \Delta} \lambda_i(x)$$

III. PRIME L -FUZZY IDEALS

Let us recall from [6] that a proper ideal P of an ADL A is said to be prime if for any $x, y \in A$, $x \wedge y \in P$ implies that $x \in P$ or $y \in P$; (equivalently, for any ideals I and J of A , $I \cap J \subseteq P \implies I \subseteq P$ or $J \subseteq P$.)

The following definition is analogous to that of a prime ideal of A . Here after A stands for an ADL with a maximal element. An L -fuzzy ideal λ of A is called proper if $\lambda(x) \neq 1$ for some $x \in A$.

Definition 3.1: A proper L -fuzzy ideal λ of A is called a prime L -fuzzy ideal if for any L -fuzzy ideals ν and μ of A , $\nu \wedge \mu \leq \lambda$ implies either $\nu \leq \lambda$ or $\mu \leq \lambda$.

An element $x \neq 1$ in L is called prime if for any $a, b \in L$ $a \wedge b \leq x$ implies either $a \leq x$ or $b \leq x$.

Now, we determine all prime L -fuzzy ideals of A by establishing a correspondence between prime L -fuzzy ideals and pairs (I, α) , where I is a prime ideal of A and α is a prime element in L . First, we recall from [7] that for any ideal I of A and $\alpha \in L$, the L -fuzzy ideal α_I of A defined by

$$\alpha_I(x) = \begin{cases} 1 & \text{if } x \in I \\ \alpha & \text{if } x \notin I. \end{cases}$$

and that α_I is called the α -level L -fuzzy ideal correspondence to I .

Theorem 3.2: Let I be an ideal of an ADL A and $\alpha \in L$. Then α_I is a prime L -fuzzy ideal of A if and only if I is a prime ideal of A and α is a prime element in L .

Proof: Suppose that α_I is a prime L -fuzzy ideal of A . Since α_I is proper, $\alpha_I(x) \neq 1$, for some $x \in A$. Therefore $x \notin I$ and hence $I \subsetneq A$. If J and K are ideals of A such that $J \cap K \subseteq I$. Then $\alpha_J \wedge \alpha_K = \alpha_{J \cap K} \leq \alpha_I$ and hence $\alpha_J \leq \alpha_I$ or $\alpha_K \leq \alpha_I$, so

that $J \subseteq I$ or $K \subseteq I$. Therefore, I is a prime ideal of A . Also, for any $\gamma, \beta \in L$,

$$\begin{aligned} \gamma \wedge \beta \leq \alpha &\implies (\gamma \wedge \beta)_I \leq \alpha_I \\ &\implies \gamma_I \wedge \beta_I \leq \alpha_I \\ &\implies \gamma_I \leq \alpha_I \text{ or } \beta_I \leq \alpha_I \\ &\implies \gamma \leq \alpha \text{ or } \beta \leq \alpha. \end{aligned}$$

Therefore, α is a prime element in L .

Conversely, suppose that I is a prime ideal of A and α is a prime element in L . Since I is proper and $\alpha \neq 1$, α_I is clearly a proper L -fuzzy ideal of A . Let λ and μ be any L -fuzzy ideals of A such that $\lambda \not\leq \alpha_I$ and $\mu \not\leq \alpha_I$. Then there exists $x, y \in A$ such that $\lambda(x) \not\leq \alpha_I(x)$ and $\mu(y) \not\leq \alpha_I(y)$. This implies that $\alpha_I(x) = \alpha = \alpha_I(y)$ (otherwise, $\alpha_I(x) = 1 \geq \lambda(x)$ and $\alpha_I(y) = 1 \geq \mu(y)$) and hence $x \notin I$ and $y \notin I$. Since I is a prime ideal, $x \wedge y \notin I$. Also, since α is prime and $\lambda(x) \not\leq \alpha$ and $\mu(y) \not\leq \alpha$, we have that $\lambda(x) \wedge \mu(y) \not\leq \alpha$.

Now, $(\lambda \wedge \mu)(x \wedge y) = \lambda(x \wedge y) \wedge \mu(x \wedge y) \geq \lambda(x) \wedge \mu(y)$ (since λ and μ are antitones) and hence $(\lambda \wedge \mu)(x \wedge y) \not\leq \alpha = \alpha_I(x \wedge y)$ so that, $(\lambda \wedge \mu) \not\leq \alpha_I$. Hence, α_I is a prime L -fuzzy ideal of A . ■

Theorem 3.3: A proper L -fuzzy ideal λ of A is prime if and only if the following are satisfied.

- (1) λ is two valued
- (2) $\lambda(m)$ is a prime element in L , for any maximal element m in A
- (3) λ_1 is a prime ideal of A .

Proof: Suppose that λ is a prime L -fuzzy ideal of A .

(1): Suppose λ assumes more than two values. Then there exists $x, y \in A$ and $\alpha \neq \beta \in L - \{1\}$ such that $\lambda(x) = \alpha$, $\lambda(y) = \beta$ and $\lambda(0) = 1$. Now, define L -fuzzy subsets ν and μ of A as follows:

$$\nu(z) = \begin{cases} 1 & \text{if } z \in [x] \\ 0 & \text{if } z \notin [x] \end{cases} \quad \text{and} \quad \mu(z) = \begin{cases} 1 & \text{if } z = 0 \\ \alpha & \text{if } z \neq 0. \end{cases}$$

Then, clearly $\nu = 0_{[x]}$ and $\mu = \alpha_{\{0\}}$ and hence ν and μ are L -fuzzy ideals. Also, for

$$z = 0 \implies (\nu \wedge \mu)(0) = \nu(0) \wedge \mu(0) = 1 \wedge 1 = 1 = \lambda(0).$$

$0 \neq z \in [x] \implies \nu(z) \wedge \mu(z) = 1 \wedge \alpha = \alpha = \lambda(x) \leq \lambda(z)$ (since λ is an antitone and $z \wedge x \leq x$, we have

$$\lambda(x) \leq \lambda(z \wedge x) = \lambda(x \wedge z) = \lambda(z))$$

and $z \notin [x] \implies \nu(z) \wedge \mu(z) = 0 \wedge \alpha = 0 \leq \lambda(z)$. Therefore, $\nu \wedge \mu \leq \lambda$. Since λ is prime, $\nu \leq \lambda$ or $\mu \leq \lambda$. But $\nu \not\leq \lambda$ (since $\nu(x) = 1, \lambda(x) = \alpha$ and $1 \neq \alpha$).

Therefore, $\mu \leq \lambda$. In particular, $\mu(y) \leq \lambda(y) \neq \lambda(0)$, we get that $y \neq 0$ and $\alpha = \mu(y) = \beta$, which is a contradiction.

(2): Let m be a maximal element in A . Since λ is proper, $\lambda(x) \neq 1$, for some $x \in A$ and hence $\lambda(m) \neq \lambda(0) = 1$

$$\left(\lambda(m) = 1 \implies \lambda(m \vee x) = 1 \implies \lambda(m) \wedge \lambda(x) = 1 \implies \lambda(x) = 1. \right)$$

Let α and $\beta \in L$ such that $\alpha \wedge \beta \leq \lambda(m)$. Define ν and μ of A as:

$$\nu(x) = \begin{cases} 1 & \text{if } x = 0 \\ \alpha & \text{if } x \neq 0 \end{cases} \quad \text{and} \quad \mu(x) = \begin{cases} 1 & \text{if } x = 0 \\ \beta & \text{if } x \neq 0. \end{cases}$$

Then, it can be easily proved that ν and μ are L -fuzzy ideals of A and $\nu \wedge \mu \leq \lambda$. Since λ is prime, $\nu \leq \lambda$ or $\mu \leq \lambda$, in particular, $\nu(m) \leq \lambda(m)$ or $\mu(m) \leq \lambda(m)$. Therefore, $\alpha \leq$

$\lambda(m)$ or $\beta \leq \lambda(m)$ and hence $\lambda(m)$ is prime.
 (3): Let $I = \{x \in A : \lambda(x) = 1\}$. Clearly, I is a proper ideal of A , since λ is proper. Let α be the other value of λ . Then

$$\lambda(x) = \begin{cases} 1 & \text{if } x \in I \\ \alpha & \text{if } x \notin I \end{cases}$$

and hence $\lambda = \alpha_I$. By theorem 3.2, I is prime. Conversely suppose that λ is an L -fuzzy ideal of A satisfying the conditions (1),(2) and (3). By (1), there exists $\alpha (\neq 1) \in L$ such that $\lambda(x) = \alpha$, for each $x \in A - \{0\}$. Then for any maximal element m of A , $\lambda(m) = \alpha$. By (2), α is prime. Let $I = \{x \in A : \lambda(x) = 1\}$. Then I is a prime ideal of A (by (3)). Therefore, $\lambda = \alpha_I$ and hence λ is a prime L -fuzzy ideal of A (by theorem 3.2). ■

The results 3.2 and 3.3 yield the following.
Theorem 3.4: Let λ be an L -fuzzy subset of A . Then λ is a prime L -fuzzy ideal of A if and only if there exists a prime ideal P of A and a prime element α in L such that $\lambda = \alpha_P$.

IV. L-FUZZY PRIME IDEALS

In this section, we introduce the notion of an L -fuzzy prime ideal which is weaker than that of a prime L -fuzzy ideal.

Definition 4.1: A proper L -fuzzy ideal λ of A is called an L -fuzzy prime ideal of A if for any $x, y \in A$, $\lambda(x \wedge y) = \lambda(x)$ or $\lambda(y)$.

The following theorem gives a characterization of an L -fuzzy prime ideal.

Theorem 4.2: Let λ be a proper L -fuzzy ideal of A . Then the following are equivalent to each other.

- (1) for each $\alpha \in L$, $\lambda_\alpha = A$ or λ_α is a prime ideal of A
- (2) λ is an L -fuzzy prime ideal of A
- (3) for any $x, y \in A$, $\lambda(x \wedge y) \leq \lambda(x) \vee \lambda(y)$ and either $\lambda(x) \leq \lambda(y)$ or $\lambda(y) \leq \lambda(x)$.

Proof: (1) \Rightarrow (2) : Let $x, y \in A$ and $\alpha = \lambda(x \wedge y)$. Then $x \wedge y \in \lambda_\alpha$ and hence $x \in \lambda_\alpha$ or $y \in \lambda_\alpha$.

$$\begin{aligned} x \in \lambda_\alpha &\Rightarrow \lambda(x \wedge y) = \alpha \leq \lambda(x) \leq \lambda(x \wedge y) \\ &\Rightarrow \lambda(x \wedge y) = \lambda(x) \end{aligned}$$

Similarly, $y \in \lambda_\alpha \Rightarrow \lambda(x \wedge y) = \lambda(y)$.
 (2) \Rightarrow (3) : Let $x, y \in A$. Then, $\lambda(x \wedge y) = \lambda(x)$ or $\lambda(y)$.

$\lambda(x \wedge y) = \lambda(x) \Rightarrow \lambda(x \wedge y) = \lambda(x) \leq \lambda(x) \vee \lambda(y)$ and $\lambda(y) \leq \lambda(x \wedge y) = \lambda(x)$.

Similarly, $\lambda(x \wedge y) = \lambda(y) \Rightarrow \lambda(x \wedge y) \leq \lambda(x) \vee \lambda(y)$ and $\lambda(x) \leq \lambda(y)$.

(3) \Rightarrow (1) : Let $\alpha \in L$ be fixed. If $\lambda_\alpha \neq A$, then λ_α is a proper ideal of A . Also, for any $x, y \in A$,

$$\begin{aligned} x \wedge y \in \lambda_\alpha &\Rightarrow \alpha \leq \lambda(x \wedge y) \leq \lambda(x) \vee \lambda(y) = \lambda(x) \text{ or } \lambda(y) \\ &\Rightarrow \alpha \leq \lambda(x) \text{ or } \alpha \leq \lambda(y) \\ &\Rightarrow x \in \lambda_\alpha \text{ or } y \in \lambda_\alpha \end{aligned}$$

Therefore, λ_α is prime. ■

Theorem 4.3: A prime L -fuzzy ideal of A is an L -fuzzy prime ideal of A .

Proof: Let λ be a prime L -fuzzy ideal of A . Then $\lambda = \alpha_I$ for some prime ideal P of A and α a prime element in L . Since $\alpha < 1$, λ is a proper. Let $x, y \in A$. Then

$$\begin{aligned} x \wedge y \in I &\Rightarrow \lambda(x \wedge y) = 1 \text{ and } x \in I \text{ or } y \in I \\ &\Rightarrow \lambda(x \wedge y) = 1 = \lambda(x) \text{ or } \lambda(y) \end{aligned}$$

$$\begin{aligned} \text{and } x \wedge y \notin I &\Rightarrow x \notin I \text{ and } y \notin I \\ &\Rightarrow \lambda(x \wedge y) = \alpha = \lambda(x) = \lambda(y) \end{aligned}$$

Therefore, λ is an L -fuzzy prime ideal of A . ■

The converse of the above theorem is not true; for consider the given example below.

Example 4.4: Let $A = \{0, a, b, c\}$, $L = \{0, t, 1\}$ with $0 < t < 1$ and let \vee and \wedge be binary operations on A defined by

\vee	0	a	b	c
0	0	a	b	c
a	a	a	a	a
b	b	b	b	b
c	c	a	b	c

\wedge	0	a	b	c
0	0	0	0	0
a	0	a	b	c
b	0	a	b	c
c	0	c	c	c

Then, $(A, \wedge, \vee, 0)$ is an ADL. Now define $\lambda : A \rightarrow L$ by $\lambda(0) = 1$, $\lambda(a) = \lambda(b) = 0$ and $\lambda(c) = t$. Therefore, $\lambda_0 = A$, $\lambda_t = \{0, c\}$ and $\lambda_1 = \{0\}$ are prime ideals of A . Therefore, λ is an L -fuzzy prime ideal of A , while λ is not a prime L -fuzzy ideal of A , since λ is not exactly two valued.

Finally, in this section we slightly generalize α -level fuzzy ideals of A and identify general prime ideals of A with L -fuzzy prime ideals of A .

Theorem 4.5: Let I a proper ideal of A and $\alpha, \beta \in L$. Let $\langle \alpha, \beta \rangle_I$ be an L -fuzzy subset of A defined by

$$\langle \alpha, \beta \rangle_I(x) = \begin{cases} 1 & \text{if } x = 0 \\ \alpha & \text{if } 0 \neq x \in I \\ \beta & \text{if } x \notin I. \end{cases}$$

Then, $\langle \alpha, \beta \rangle_I$ is an L -fuzzy ideal of A if and only if $\beta \leq \alpha$ and, in this case $\langle \alpha, \beta \rangle_I$ is proper if and only if $\beta < 1$.

I is a prime ideal of A if and only if χ_I is an L -fuzzy prime ideal of A

(3) Suppose that 0 be a prime element in L . Then, I is a prime ideal of A if and only if $\langle \alpha, \beta \rangle_I$ is an L -fuzzy prime ideal of A for all $1 \neq \beta \leq \alpha$ in L .

Proof: (1) and (2) are straight forward and simple verifications.

(3): Suppose that I is a prime ideal of A and $1 \neq \beta \leq \alpha$ in L . Let $x, y \in I$. Then,

$$\begin{aligned} x \wedge y = 0 &\Rightarrow x = 0 \text{ or } y = 0 \\ &\Rightarrow \langle \alpha, \beta \rangle_I(x \wedge y) = 1 = \langle \alpha, \beta \rangle_I(x) \text{ or } \langle \alpha, \beta \rangle_I(y) \\ 0 \neq x \wedge y \in I &\Rightarrow 0 \neq x \in I \text{ or } 0 \neq y \in I \\ &\Rightarrow \langle \alpha, \beta \rangle_I(x \wedge y) = \alpha = \langle \alpha, \beta \rangle_I(x) \text{ or } \langle \alpha, \beta \rangle_I(y) \end{aligned}$$

and $x \wedge y \notin I \Rightarrow x \notin I$ and $y \notin I$
 $\Rightarrow \langle \alpha, \beta \rangle_I(x \wedge y) = \beta = \langle \alpha, \beta \rangle_I(x) = \langle \alpha, \beta \rangle_I(y)$.
 Converse follows from the fact that $\chi_I = \langle 1, 0 \rangle_I$. ■

V. MINIMAL PRIME L-FUZZY IDEALS

Let us recall from [?] that a prime ideal P an ADL of A containing an ideal I is said to be a minimal prime ideal belonging to I if there is no prime ideal of A containing I and properly contained in P .

Definition 5.1: Let λ be a prime L -fuzzy ideal of A . Then λ is said to be minimal if λ is a minimal member in the set of all prime L -fuzzy ideals of A under the point-wise partial ordering. A minimal prime L -fuzzy ideal belonging to $\chi_{\{0\}}$ is simply called a minimal prime L -fuzzy ideal.

In this section, we characterize all minimal prime L -fuzzy ideals of A in terms of minimal prime ideals of A and minimal prime elements of L .

As usual, by a minimal prime element of L we mean a minimal element in the poset of all prime elements of L .

Now we have the following:

Theorem 5.2: Let λ be an L -fuzzy ideal of A . Then λ is a minimal prime L -fuzzy ideal of A if and only if $\lambda = \alpha_I$, for some minimal prime ideal I of A and a minimal prime element α in L .

Proof: Suppose that $\lambda = \alpha_I$ for some minimal prime ideal I of A and minimal prime element in L . Then by theorem 3.4, λ is prime L -fuzzy ideal of A . Let μ be a prime L -fuzzy ideal of A and $\mu \leq \lambda$. Then by theorem 3.4, $\mu = \beta_J$ for some prime ideal J of A and a prime element β in L . Therefore, $\beta_J \leq \alpha_I$. This implies that, $\beta \leq \alpha$ and $J \subseteq I$. By using the minimality of I and α , we get that $\beta = \alpha$ and $J = I$. Therefore, $\mu = \lambda$ and hence λ is a minimal prime L -fuzzy ideal of A .

Conversely suppose that λ is a minimal prime L -fuzzy ideal of A . Then by theorem 3.4, there exists a prime ideal I of A and a prime element α in L such that

$\lambda = \alpha_I$. Let J be a prime ideal of A such that $J \subseteq I$. Then $\alpha_J \leq \alpha_I$, by the minimality of λ , $\alpha_J = \alpha_I$. Therefore, $J = I$ and hence I is minimal prime ideal of A . Let β be a prime element in L and $\beta \leq \alpha$. Then $\beta_I \leq \alpha_I$. This implies, $\beta_I = \alpha_I$ and hence $\beta = \alpha$. Thus α is a minimal prime element in L . ■

If the smallest element 0 in L is prime, then 0 will be the only minimal prime element in L . Note that $\chi_P = 0_P$, for any ideal P of A .

The following is a simple verification.

Theorem 5.3: Let 0 be a prime element in L . Then an L -fuzzy ideal λ of A is a minimal prime L -fuzzy ideal of A if and only if $\lambda = \chi_P$, for some minimal prime ideal P of A . More over, $P \mapsto \chi_P$ is a bijection of the set of minimal prime ideals of A onto the set of minimal prime L -fuzzy ideals of A .

VI. L-FUZZY MINIMAL PRIME IDEALS

By an L -fuzzy minimal prime ideal of A we mean, as usual, a minimal element in the set of all L -fuzzy prime ideals of A under the point-wise partial ordering. In this section, we characterize all L -fuzzy minimal prime ideals of A in terms of their α -cuts.

Theorem 6.1: (1) If λ is an L -fuzzy prime ideal of A , then $\lambda_1 = \{x \in A : \lambda(x) = 1\}$ is a prime ideal of A .

(2) Let λ be an L -fuzzy prime ideal of A . If λ is an L -fuzzy minimal prime ideal of A , then λ_1 is a minimal prime ideal of A .


Proof: (1) Let λ be an L -fuzzy prime ideal of A . Then λ_1 is a proper ideal of A since λ is proper. Let $x, y \in A$. Then, $x \wedge y \in \lambda_1 \Rightarrow \lambda(x \wedge y) = 1$

$$\Rightarrow 1 = \lambda(x \wedge y) = \lambda(x) \text{ or } \lambda(y) \text{ (by 4.1)}$$

$$\Rightarrow x \in \lambda_1 \text{ or } y \in \lambda_1.$$

Thus, λ_1 is a prime ideal of A .

The converse is not true. For, consider the lattice $A = \{0, a, b, c, 1\}$ represented by the Hasse diagram is given below.



Define $\lambda : A \rightarrow [0, 1]$ by $\lambda(0) = 1, \lambda(c) = 0.75, \lambda(b) = 0.5$ and $\lambda(a) = \lambda(1) = 0$. Then, $\lambda_1 = \{0\}$ which is a prime ideal of A , while λ is not an L -fuzzy prime ideal of A , since $\lambda(a \wedge b) = \lambda(c) = 0.75 \neq \lambda(a)$ and $\lambda(b)$.

(2) Suppose that λ is an L -fuzzy minimal prime ideal of A . Let Q be a prime ideal of A and $Q \subset \lambda_1$. Then χ_Q is an L -fuzzy prime ideal of A and $\chi_Q \not\leq \lambda$. This implies that λ is not an L -fuzzy minimal prime ideal of A , which is a contradiction. Thus λ_1 is a minimal prime ideal of A .

The converse is not true; for in the above example, if $\lambda(0) = 1$ and $\lambda(x) = 0.5$ for all $x \neq 0$, then it can be easily checked that λ is an L -fuzzy prime ideal of A and $\lambda_\alpha = A$ if $0 \leq \alpha \leq 0.5$ and $\lambda_\alpha = \{0\}$ if $0.5 < \alpha \leq 1$. In particular, λ_1 is a minimal prime ideal of A . But, λ is not an L -fuzzy minimal prime ideal of A , since if we define $\mu(0) = 1$ and $\mu(x) = 0.25$ for all $x \neq 0$, then μ is an L -fuzzy prime ideal of A and $\mu \not\leq \lambda$. ■

The following theorem is a characterization of L -fuzzy minimal prime ideals of A .

Theorem 6.2: Let λ be an L -fuzzy prime ideal of A and 0 be a prime element in L . Then λ is an L -fuzzy minimal prime ideal of A if and only if λ_α is a minimal prime ideal of A , for all $\alpha \in L$.

Proof: Suppose λ is an L -fuzzy minimal prime ideal of A and λ_α is not minimal prime ideal of A , for some $0 < \alpha < 1$ in L . Then there exists a prime ideal Q of A such that $Q \subset \lambda_\alpha$. Define $\mu : A \rightarrow L$ by

$$\mu(x) = \begin{cases} 1 & \text{if } x = 0 \\ \alpha & \text{if } 0 \neq x \in Q \\ 0 & \text{if } x \notin Q. \end{cases}$$


Then, clearly $\mu = (\alpha, 0)_Q$ and hence μ is an L -fuzzy prime ideal of A (by theorem 4.5 (3)). Also, $\mu \leq \lambda$. Since $Q \subset \lambda_\alpha$, there exists $y \in \lambda_\alpha$ such that $y \notin Q$. Therefore, $\mu(y) = 0 <$

$\alpha \leq \lambda(y)$. Therefore, $\mu \not\leq \lambda$, which is a contradiction. Thus for each $\alpha \in L$, λ_α is a minimal prime ideal of A .

Conversely, suppose for each $\alpha \in L$, λ_α is a minimal prime ideal of A . Let μ be an L -fuzzy prime ideal of A such that $\mu \leq \lambda$. Then for each $\alpha \in L$, $\mu_\alpha \subseteq \lambda_\alpha$. By the minimality of λ_α , we have $\mu_\alpha = \lambda_\alpha$ and hence $\mu = \lambda$. Therefore λ is an L -fuzzy minimal prime ideal of A . ■

Remark 6.3: If λ is an L -fuzzy minimal prime ideal of A , the each α -cut of λ need not be minimal prime ideal of A .

For, consider the example given in the following. Let $A = \{0, a, b, c, 1\}$ be the lattice represented by the Hasse diagram is given below.



Define $\lambda : A \rightarrow [0, 1]$ by $\lambda(0) = \lambda(a) = 1$, $\lambda(b) = \lambda(c) = 0.5$ and $\lambda(1) = 0$. It can be easily verified that, λ is an L -fuzzy prime ideal of A and for any $t \in [0, 0.5]$, $\lambda_t = \{0, a, b, c\}$ is a prime ideal of A but not minimal.

REFERENCES

- [1] L. Zadeh, "Information and control," *Fuzzy sets*, vol. 8, no. 3, pp. 338–353, 1965.
- [2] J. Goguen, "L-fuzzy sets," *Journal of mathematical analysis and applications*, vol. 18, no. 1, pp. 145–174, 1967.
- [3] W.-j. Liu, "Fuzzy invariant subgroups and fuzzy ideals," *Fuzzy sets and systems*, vol. 8, no. 2, pp. 133–139, 1982.
- [4] T. Mukherjee and M. Sen, "On fuzzy ideals of a ring i," *Fuzzy Sets and systems*, vol. 21, no. 1, pp. 99–104, 1987.
- [5] U. Swamy and K. Swamy, "Fuzzy prime ideals of rings," *Journal of Mathematical Analysis and Applications*, vol. 134, no. 1, pp. 94–103, 1988.
- [6] U. Swamy and G. Rao, "Almost distributive lattices," *Journal of the Australian Mathematical Society*, vol. 31, no. 1, pp. 77–91, 1981.
- [7] U. Swamy, C. S. S. Raj, and N. Teshale, "Fuzzy ideals of almost distributive lattices," *Annals of Fuzzy Mathematics and Informatics*, vol. 14, no. 4, pp. 371–379, 2017.